

Exercise sheet 8  
Theoretical Physics 3 : QM SS2018  
Lecturer : Prof. M. Vanderhaeghen

13.06.2018

**Exercise 0.**

How much time did it take to complete the task?

**Exercise 1. Infinite spherical well (30 points)**

Consider a particle in an infinite well 3D potential of radius  $a$ ,

$$V(r) = \begin{cases} 0 & r < a, \\ +\infty & r \geq a. \end{cases}$$

a) (15 p.) Show that the solution of the Schrödinger equation is

$$\Psi_{nlm}(r, \theta, \phi) \propto j_l\left(\beta_{nl}\frac{r}{a}\right) Y_{lm}(\theta, \phi),$$

and  $j_l(x)$  is the spherical Bessel functions of order  $l$  which is defined as:

$$j_l(x) \equiv (-x)^l \left(\frac{1}{x} \frac{d}{dx}\right)^l \frac{\sin x}{x}.$$

$j_l(x)$  is the nonsingular at zero solution of the differential equation

$$x^2 \frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} + [x^2 - n(n+1)] y = 0.$$

$\beta_{nl}$  is  $n$ th zero of the a spherical Bessel functions of order  $l$ :  $j_l(\beta_{nl}) = 0$ .

b) (15 p.) The spherical Bessel functions is a particular case of the Bessel functions  $J_\alpha(x)$  defined as:

$$J_\alpha = \sum_{n=0}^{\infty} \frac{(-1)^n}{n! \Gamma(n + \alpha + 1)} \left(\frac{x}{2}\right)^{2n + \alpha},$$

for  $\alpha$  being half-integer, so  $J_{n+1/2} = \sqrt{\frac{2x}{\pi}} j_n(x)$ .

Using the definition of the Bessel functions, compute  $J_{1/2}$  and  $J_{3/2}$  and check that, indeed, the relation between  $J_{l+1/2}$  and  $j_l$  is correct.

*Hint:* Prove  $n!(1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n+1))2^n = (2n+1)!$ .

*Math hints:*

$$\sin(x) = \sum_{m=0}^{\infty} (-1)^m \frac{x^{2m+1}}{(2m+1)!}$$
$$\Gamma(m+1/2) = \frac{1 \cdot 3 \cdot 5 \cdot (2m-1)}{2^m} \sqrt{\pi}$$

## Exercise 2. Hydrogen atom (20 + 10 points)

The normalized hydrogen wave functions are:

$$\psi_{nlm}(r, \theta, \phi) = \frac{2}{n^2} \sqrt{\frac{(n-l-1)!}{[a(n+l)!]^3}} e^{-\frac{r}{na}} \left(\frac{2r}{na}\right)^l L_{n-l-1}^{2l+1}\left(\frac{2r}{na}\right) Y_l^m(\theta, \phi),$$

where  $L_{q-p}^p(x)$  are the associated Laguerre polynomials and  $Y_l^m(\theta, \phi)$  are the spherical harmonics.

- (5 p.) Consider the electron is in the state  $\psi_{nlm}(r, \theta, \phi)$ . What is the probability  $P_{nl}(r)$  to find it somewhere?
- (15 p.) Check explicitly that  $P_{nl}(r)$  is correctly normalized to unity for  $n = 3$ .  
*Hint:* Use  $\int_0^\infty dx e^{-x} x^n = n!$ .
- (10 p.) (*Bonus*) Show that  $\int_0^\infty dx e^{-x} x^n = n!$ .

## Exercise 3. 2D quantum harmonic oscillator. (50 + 20 points)

- (10 p.) Assuming solutions for the one-dimensional case are already known, solve the two-dimensional isotropic quantum harmonic oscillator problem in the Cartesian coordinates:

$$-\frac{\hbar^2}{2m} \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \psi(x, y) + \frac{m\omega^2}{2} (x^2 + y^2) \psi(x, y) = E\psi(x, y).$$

*Hint:* Use the method of separation of variables:  $\psi(x, y) = X(x)Y(y)$ . Then write down separate equations on  $X(x)$  and  $Y(y)$ .

- (5 p.) Write down the energy spectrum. What is the degree of degeneracy of the energy levels?
- (10 p.) Show that the Laplace operator in two dimensions in the polar coordinates takes the form

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2}.$$

- (10 p.) Write down the angular momentum operator  $\hat{L}_z = \hat{x}\hat{p}_y - \hat{y}\hat{p}_x$  in the polar coordinates and show that  $[\hat{H}, \hat{L}_z] = 0$ .
- (10 p.) Consider the two-dimensional isotropic harmonic oscillator in polar coordinates. Separate variables  $\psi(r, \phi) = v(r)u(\phi)$  and get equations on  $v(r)$  and  $u(\phi)$ .

The equation on  $v(r)$  can be eventually transformed into the one for the generalised Laguerre polynomials  $L_{n_r}^{|M|+1}\left(\frac{m\omega}{\hbar}r^2\right)$ . Then one obtains the final solution

$$\psi_{n_r M}(r, \phi) = C_{n_r M} r^{|M|} e^{-\frac{m\omega}{2\hbar}r^2} L_{n_r}^{|M|+1}\left(\frac{m\omega}{\hbar}r^2\right) e^{iM\phi}$$

with the spectrum

$$E = \hbar\omega(|M| + 1 + 2n_r), \quad n_r = 0, 1, 2, \dots, \quad M = 0, \pm 1, \pm 2, \dots,$$

where  $M$  is the quantum number corresponding to  $\hat{L}_z$ .

- (5 p.) Find eigenvalues and eigenfunctions of  $\hat{L}_z$  in polar coordinates. Show that, indeed, the complete and orthonormal set of eigenfunctions is common to both  $\hat{H}$  and  $\hat{L}_z$ .
- (20 p.) (*Bonus*) Find the ground state solution of the Schrödinger equation ( $n_r = 0, M = 0$ ) in polar coordinates.

*Hint:* put  $E = \hbar\omega$ ,  $u''(\phi) = 0$  and substitute  $v(r) = e^{-\frac{m\omega}{2\hbar}r^2} F(r)$ .