

A glimpse of the H dibaryon from a lattice QCD perspective

Hartmut Wittig

PRISMA⁺ Cluster of Excellence, Institute for Nuclear Physics and Helmholtz Institute Mainz

58th International Winter Meeting on Nuclear Physics

Bormio, Italy

20 – 24 January 2020

Introduction – The *H* dibaryon

VOLUME 38, NUMBER 5

PHYSICAL REVIEW LETTERS

31 JANUARY 1977

Perhaps a Stable Dihyperon*

R. L. Jaffe†

*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305, and Department of Physics
and Laboratory of Nuclear Science,‡ Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

(Received 1 November 1976)

In the quark bag model, the same gluon-exchange forces which make the proton lighter than the $\Delta(1236)$ bind six quarks to form a stable, flavor-singlet (with strangeness of -2) $J^P = 0^+$ dihyperon (H) at 2150 MeV. Another isosinglet dihyperon (H^*) with $J^P = 1^+$ at 2335 MeV should appear as a bump in $\Lambda\Lambda$ invariant-mass plots. Production and decay systematics of the H are discussed.

- * MIT bag model predicts dihyperon state (H) with

$$I = 0, S = -2, J^P = 0^+$$

and a mass of $m_H = 2150 \text{ MeV}$

- * H dibaryon must decay weakly

Experimental Searches

VOLUME 87, NUMBER 21

PHYSICAL REVIEW LETTERS


19 NOVEMBER 2001

Observation of a $_{\Lambda\Lambda}^6\text{He}$ Double Hypernucleus

(E373@KEK):

A double-hyperfragment event has been found in a hybrid-emulsion experiment. It is identified uniquely as the sequential decay of $_{\Lambda\Lambda}^6\text{He}$ emitted from a Ξ^- hyperon nuclear capture at rest. The mass of $_{\Lambda\Lambda}^6\text{He}$ and the Λ - Λ interaction energy $\Delta B_{\Lambda\Lambda}$ have been measured for the first time devoid of the ambiguities due to the possibilities of excited states. The value of $\Delta B_{\Lambda\Lambda}$ is $1.01 \pm 0.20^{+0.18}_{-0.11}$ MeV. This demonstrates that the Λ - Λ interaction is weakly attractive.

“Nagara” event



Observation of a $_{\Lambda\Lambda}^6\text{He}$ double-hypernucleus

Binding energy:

$$B_{\Lambda\Lambda} = 7.25 \pm 0.19 (^{+0.18}_{-0.11}) \text{ MeV}$$

Interpreted as sequential weak decay of $_{\Lambda\Lambda}^6\text{He}$

$$m_H > 2m_\Lambda - B_{\Lambda\Lambda} = 2223.7 \text{ MeV} \quad @ 90\% \text{ CL}$$

The H dibaryon as a dark matter candidate

- * $udsuds$ bound state as dark matter candidate:

[G.R. Farrar, A. Strumia et al.,...]

$$m_H < 2(m_p + m_e) = 1877.6 \text{ MeV} \quad \Rightarrow \quad H \text{ dibaryon absolutely stable}$$

$$m_H > 2(m_p + \text{B.E.}) = 1860 \text{ MeV} \quad \Rightarrow \quad \text{Nuclei absolutely stable}$$



- * Recall: $2m_\Lambda = 2230 \text{ MeV}$

$$m_H = 2150 \text{ MeV} \quad (\text{Jaffe's bag model estimate})$$

- * Scenario requires very large binding energy of $\approx 360 \text{ MeV}$

Current Status

- * H dibaryon not firmly established experimentally
- * Is a bound H dibaryon a consequence of QCD?
- * Try “ab initio” technique: Lattice QCD



“Clover” @ Mainz

Beyond Perturbation Theory: Lattice QCD

- * Non-perturbative treatment; regularised Euclidean functional integrals

Lattice spacing: $a, \quad x_\mu = n_\mu a, \quad a^{-1} = \Lambda_{\text{UV}}$

Finite volume: $L^3 \cdot T, \quad N_s = L/a, \quad N_t = T/a$

$$\langle \Omega \rangle = \frac{1}{Z} \int \prod_{x,\mu} dU_\mu(x) \Omega \prod_{f=u,d,s,\dots} \det(\not{D}^{\text{lat}} + m_f) e^{-S_G[U]}$$

- * Stochastic evaluation of $\langle \Omega \rangle$ via Markov process

Strong growth of numerical cost near physical m_u, m_d

- * Pion mass, i.e. lightest mass in pseudoscalar channel:

$$\begin{array}{ccc} \approx 500 \text{ MeV} & \longrightarrow & \approx 130 \text{ MeV} \\ (2001) & & (\gtrsim 2015) \end{array}$$

Hadron spectrum in Lattice QCD

- * Spectral information contained in correlation functions

$$\sum_{x,y} e^{ip \cdot (y-x)} \langle O_{\text{had}}(y) O_{\text{had}}^\dagger(x) \rangle = \sum_n w_n(\mathbf{p}) e^{-E_n(\mathbf{p})(y_0-x_0)}$$
$$\xrightarrow{(y_0-x_0) \rightarrow \infty} w_1(\mathbf{p}) e^{-E_1(\mathbf{p})(y_0-x_0)}$$

- * $O_{\text{had}}(x)$: **interpolating operator**
 - projects on all states with the same quantum numbers

Nucleon: $O_N = \epsilon_{abc} (u^a C \gamma_5 d^b) u^c$

- * Ground state dominates at large Euclidean times: $y_0 - x_0 \rightarrow \infty$
- * Excited states are **sub-leading** contributions

The H Dibaryon in Lattice QCD

Flavour structure

- * H dibaryon lies in the 1-dimensional irrep. of $SU(3)_{\text{flavour}}$

- * Flavour structure of two octet baryons:

$$\mathbf{8} \otimes \mathbf{8} = (\mathbf{1} \oplus \mathbf{8} \oplus \mathbf{27})_S \oplus (\mathbf{8} \oplus \mathbf{10} \oplus \overline{\mathbf{10}})_A$$

- * Upon $SU(3)$ -symmetry breaking, $\mathbf{8}$ and $\mathbf{27}$ mix with singlet
- * Singlet, octet and $\mathbf{27}$ plet interpolators constructed from linear combinations of $\Lambda\Lambda$, $\Sigma\Sigma$ and $N\Xi$ operators

Other interesting dibaryons

- * Dineutron lies in $\mathbf{27}$ irrep.
- * Deuteron lies in $\overline{\mathbf{10}}$ irrep. with $J^P = 1^+$

Interpolating operators

- * Hexaquark operators (inspired by Jaffe's original bag model calculation):

$$[rstuvw] = \epsilon_{ijk} \epsilon_{lmn} (s^a C \gamma_5 P_+ t^b) (v^l C \gamma_5 P_+ w^m) (r^k C \gamma_5 P_+ u^n)$$

$$H^{(1)} = \frac{1}{48} ([sudsud] - [udusds] - [dudsus])$$

$$H^{(27)} = \frac{1}{48\sqrt{3}} (3[sudsud] + [udusds] - [dudsus])$$

- * Momentum-projected two-baryon operators:

$$B_\alpha \equiv [rst]_\alpha = \epsilon_{ijk} (s^i C \gamma_5 P_+ t^j) r_\alpha^k$$

$$(BB)(\mathbf{P}; t) = \sum_{\mathbf{x}} e^{-i\mathbf{p}_1 \cdot \mathbf{x}} B_1(\mathbf{x}, t) (C \gamma_5 P_+) \sum_{\mathbf{y}} e^{-i\mathbf{p}_2 \cdot \mathbf{y}} B_2(\mathbf{y}, t), \quad \mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2$$

→ project onto $(BB)^{(1)}, (BB)^{(8)}, (BB)^{(27)}$

Correlation matrices

- * Consider set of N_{op} interpolating operators for a given hadron:

Correlation matrix: $C_{ij}(\mathbf{P}, \tau) = \langle O_i(\mathbf{P}, t) O_j(\mathbf{P}, t')^\dagger \rangle, \quad \tau = t - t'$

- * **Variational method:** solve Generalised Eigenvalue Problem (GEVP):

$$\mathbf{C}(t_1) v_n(t_1, t_0) = \lambda_n(t_1, t_0) \mathbf{C}(t_0) v_n(t_1, t_0)$$

$$w_n^\dagger(t_1, t_0) \mathbf{C}(t_1) = \lambda_n(t_1, t_0) w_n^\dagger(t_1, t_0) \mathbf{C}(t_0), \quad n = 1, \dots, N_{\text{op}}$$

- * Project on approximately diagonal correlator:

$$\Lambda_{mn}(t) = w_n^\dagger \mathbf{C}(t) v_m$$

- * Compute the effective n^{th} energy level:

$$E_n^{\text{eff}}(t) = \frac{1}{\Delta t} \ln \frac{\Lambda_{nn}(t)}{\Lambda_{nn}(t + \Delta t)}$$

HAL QCD Method

- * Obtain baryon-baryon potential from Nambu-Bethe-Salpeter amplitude computed on the lattice

$$G_4(\mathbf{r}, t - t_0) = \langle 0 | (BB)^{(\alpha)}(\mathbf{r}, t) (\overline{B}\overline{B})^{(\alpha)}(\mathbf{r}, t_0) | 0 \rangle = \phi(\mathbf{r}, t) e^{-2M(t-t_0)}$$

$(BB)^{(\alpha)}(\mathbf{r}, t)$: 2-baryon interpolating operator; flavour irrep. α

$\phi(\mathbf{r}, t)$: NBS wave function

M : single baryon mass

- * Determine potential via

$$V(r) = \frac{[-H_0 - (\partial/\partial t)] \phi(\mathbf{r}, t)}{\phi(\mathbf{r}, t)}$$

- * Solve Schrödinger equation

→ determine binding energies and scattering phase shifts

HAL QCD Method


Details of the calculation:

[Inoue et al., Phys Rev Lett 106 (2011) 162002]

$N_f = 3$, i.e. mass-degenerate u, d, s quarks

Single lattice spacing: $a = 0.121(2)$ fm

5 pion masses in the range: $m_\pi = 469 - 1171$ MeV



(statistical and systematic
errors combined)

HAL QCD Method


Details of the calculation:

[Sasaki et al., arXiv:1912.08630]

$N_f = 2 + 1$, $O(a)$ improved Wilson fermions

Single lattice spacing: $a = 0.0846$ fm; Volume: $L \approx 8.1$ fm

Near physical point: $m_\pi = 146$ MeV, $m_K = 525$ MeV



- * $\Lambda\Lambda$ interaction weakly attractive
- ⇒ No bound or resonant dihyperon near $\Lambda\Lambda$ threshold at the physical point

Needs verification using
a different methodology

The Mainz Dibaryon Project

Collaborators:


A. Francis, J.R. Green, A. Hanlon, P. Junnarkar, Ch. Miao, T.D. Rae, H.W.

Gauge ensembles provided by the CLS effort:

- * $N_f = 2$ flavours of $\mathcal{O}(a)$ improved Wilson fermions; quenched strange quark
Pion masses: $m_\pi = 450 - 1000 \text{ MeV}$ (to compare with earlier studies)
[Francis, Green, Junnarkar, Miao, Rae, HW, Phys Rev D99 (2019) 074505]
- * $N_f = 2 + 1$ flavours of $\mathcal{O}(a)$ improved Wilson fermions
Pion masses: $m_\pi = 200 - 420 \text{ MeV}$
[Hanlon, Francis, Green, Junnarkar, HW, arXiv:1810.13282]
- * SU(3)-symmetric and SU(3)-broken situations
- * Three different lattice spacings to investigate lattice artefacts

Finite-volume spectrum

- * Ensemble E1: $N_f = 2$, SU(3)-symmetric, $m_\pi \approx 960$ MeV
- * Point-to-all propagators: hexaquark operators at source, two-baryon or hexaquark operators at the sink




- * Hexaquark operators: noisier, slower convergence towards ground state

Finite-volume spectrum

- * Compute timeslice-to-all propagators
→ “**distillation**” — Laplace-Heaviside (LapH) smearing

[Pardon et al., PRD 80 (2009) 054506; Morningstar et al., PRD 83 (2011) 114505]



- * Quark propagator with smearing matrix at source and sink:




$$\mathcal{S} D^{-1} \mathcal{S}, \quad \mathcal{S}^{(t)}(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{N_{\text{LapH}}} V^{(k)}(\mathbf{x}, t) \otimes V^{(k)}(\mathbf{y}, t)^\dagger$$

$V^{(k)}$: k^{th} eigenvector of Laplacian Δ ; has support on entire timeslice

Finite-volume spectrum

- * Compute timeslice-to-all propagators
→ “**distillation**” — Laplace-Heaviside (LapH) smearing

[Pardon et al., PRD 80 (2009) 054506; Morningstar et al., PRD 83 (2011) 114505]






- * Much better statistical signal

Finite-volume spectrum

- * Compute timeslice-to-all propagators
→ “**distillation**” — Laplace-Heaviside (LapH) smearing

[Pardon et al., PRD 80 (2009) 054506; Morningstar et al., PRD 83 (2011) 114505]



- * Much better statistical signal
- * Ensemble E5: broken SU(3)-flavour symmetry ($m_\pi = 450$ MeV)

Scattering phase shifts – Lüscher method

* Scattering momentum: $p^2 = \frac{1}{4}(E^2 - \mathbf{P} \cdot \mathbf{P}) - m_\Lambda^2$



* Scattering phase shifts: $p \cot \delta(p) = \frac{2}{\gamma L \sqrt{\pi}} Z_{00}(1, q^2), \quad q = \frac{pL}{2\pi}$

[Lüscher 1990/91;
Rummukainen & Gottlieb 1995]

$$Z_{00}(1, q^2) = \frac{1}{\sqrt{4\pi}} \left\{ \sum_{q^2 \neq n^2}^{\Lambda_n} \frac{1}{q^2 - n^2} - 4\pi \Lambda_n \right\}$$


Scattering phase shifts – Lüscher method

- * Scattering momentum: $p^2 = \frac{1}{4}(E^2 - \mathbf{P} \cdot \mathbf{P}) - m_\Lambda^2$
- * Scattering phase shifts: $p \cot \delta(p) = \frac{2}{\gamma L \sqrt{\pi}} Z_{00}(1, q^2), \quad q = \frac{pL}{2\pi}$
- * Pole of the scattering amplitude: $\mathcal{A} \propto \frac{1}{p \cot \delta(p) - ip}$
- * Fit to effective range expansion: $p \cot \delta_0(p) = A + Bp^2 + \dots \stackrel{!}{=} -\sqrt{-p^2}$



Comparison with other calculations

- * **NPLQCD Collaboration:** point-to-all propagators
- * **HAL QCD Collaboration:** energy levels from lattice calculation of NBS wavefunction ($N_f = 3$)



Binding energy from FV analysis
(SU(3)-symmetric case)

$$B_{\Lambda\Lambda} = 19 \pm 10 \text{ MeV}$$


$$(m_\pi = 960 \text{ MeV})$$

(Green: SU(3)-symmetric; blue: SU(3)-broken)

[Francis et al., PRD 99 (2019) 074505;
Green et al., in prep.]

Comparison with other calculations

- * **NPLQCD Collaboration:** point-to-all propagators
- * **HAL QCD Collaboration:** energy levels from lattice calculation of NBS wavefunction ($N_f = 3$)



(Green: SU(3)-symmetric; blue: SU(3)-broken)

Binding energy from FV analysis
(SU(3)-symmetric case)

$$B_{\Lambda\Lambda} = 16.3 \pm 4.2 \text{ MeV}$$

$$(m_\pi = 960 \text{ MeV})$$

$$B_{\Lambda\Lambda} = 4.5 \pm 3.7 \text{ MeV}$$

$$(m_\pi = 450 \text{ MeV})$$

[Francis et al., PRD 99 (2019) 074505;
Green et al., in prep.]

Higher spin states

- * So far: focus on $J^P = 0^+$ and $S = -2$
- * Extend calculation to higher spins → include additional irreps.
- * Spin-1 interpolating operators:

$$B_\alpha \equiv [rst]_\alpha = \epsilon_{ijk} \left(s^i C \gamma_5 P_+ t^j \right) r_\alpha^k$$

$$(BB)_i(\mathbf{p}_1, \mathbf{p}_2) = \sum_{\mathbf{x}} e^{-i\mathbf{p}_1 \cdot \mathbf{x}} B_1(\mathbf{x}, t) (C \gamma_i P_+) \sum_{\mathbf{y}} e^{-i\mathbf{p}_2 \cdot \mathbf{y}} B_2(\mathbf{y}, t)$$

$$\Rightarrow \text{Deuteron: } (BB)_{i; T_1^+}^{(n)} = \frac{1}{N} \sum_{\mathbf{p}; p^2=n} (BB)_i(-\mathbf{p}, \mathbf{p})$$

- * Study H dibaryon and additional states in QCD with $N_f = 2 + 1$
- * Move toward physical pion mass

Gauge ensembles with $N_f = 2+1$

- * $\mathcal{O}(a)$ improved Wilson fermions — CLS effort
- * SU(3)-symmetric point: $m_\pi = m_K \approx 420 \text{ MeV}$

Label	$L^3 \times T$	$a \text{ [fm]}$	$m_\pi \text{ [MeV]}$	
U103	$24^3 \times 128$	0.0865	420	SU(3)-symmetric
H101	$32^3 \times 96$	0.0865	420	SU(3)-symmetric
B450	$32^3 \times 64$	0.0765	415	SU(3)-symmetric
U102	$24^3 \times 128$	0.0865	350	SU(3)-broken
D200	$64^3 \times 128$	0.0644	200	SU(3)-broken


- * Approach physical point along chiral trajectory defined by

$$\text{Tr } M_q = \text{const.} \Leftrightarrow \frac{1}{2}m_\pi^2 + m_K^2 \approx \text{const.}$$

- * Compute spectrum using distillation and GEVP

Preliminary results for $N_f = 2+1$


- * Singlet channel, spin-0, SU(3)-symmetric
- * Finite-volume energy levels in A_1 irrep. in different frames



U103 ensemble: $m_\pi = m_K = 420 \text{ MeV}$, $L = 2.08 \text{ fm}$, $a = 0.0865 \text{ fm}$

Preliminary results for $N_f = 2+1$


- * Singlet channel, spin-0, SU(3)-symmetric
- * Finite-volume energy levels in A_1 irrep. in different frames



H101 ensemble: $m_\pi = m_K = 420 \text{ MeV}$, $L = 2.77 \text{ fm}$, $a = 0.0865 \text{ fm}$

Preliminary results for $N_f = 2+1$


- * Singlet channel, spin-0, SU(3)-symmetric
- * Finite-volume energy levels in A_1 irrep. in different frames



B450 ensemble: $m_\pi = m_K = 415 \text{ MeV}$, $L = 2.45 \text{ fm}$, $a = 0.0765 \text{ fm}$

Preliminary results for $N_f = 2+1$


- * Singlet channel, spin-0, SU(3)-**broken**
- * Finite-volume energy levels in A_1 irrep. in different frames



U102 ensemble: $m_\pi = 350 \text{ MeV}$, $m_K = 450 \text{ MeV}$, $L = 2.08 \text{ fm}$, $a = 0.0865 \text{ fm}$

Preliminary results for $N_f = 2+1$


- * Singlet channel, spin-0, SU(3)-**broken**
- * Finite-volume energy levels in A_1 irrep. in different frames



- * Excellent resolution of excitation spectrum
- * Interpretation of the energy levels very involved

Preliminary results for $N_f = 2+1$

- * Is the deuteron bound at $m_\pi = m_K \approx 420 \text{ MeV}$?



Summary

- * **Distillation & GEVP:** powerful method to determine energy levels in a finite volume for a wide range of dibaryon channels
- * Lüscher's finite-volume quantisation condition: rigorous formalism to study hadron-hadron interactions on the lattice
- * **H dibaryon:** Binding energies extracted from Lüscher formalism significantly lower than the naïve energy difference $E_{\Lambda\Lambda} - 2m_\Lambda$
- * SU(3)-symmetric point: $B_{\Lambda\Lambda} = 5 - 20 \text{ MeV}$, $m_\pi \geq 450 \text{ MeV}$
⇒ significantly lower than Jaffe's bag model estimate

Next steps:

- * Investigate SU(3)-breaking
- * Compute binding energies as the quark masses are tuned towards the physical situation