

Exercise sheet 13
Theoretical Physics 5 : WS 2021/2022
Lecturers : Prof. M. Vanderhaeghen, Dr. I. Danilkin

24.01.2022

Exercise 1. (50 points) :
Path integral in terms of effective action

In Solid State physics, the non-relativistic motion of electrons is often considered perturbed by the effect of impurities, as it happens in typical semiconductors. In simple cases, within one-electron approximation, one can approximate this effect with a position-dependent effective mass Schrodinger equation.

Consider a one-dimensional system with position-dependent mass:

$$H = \frac{p^2}{2f(x)}.$$

- a) (5 p.) Show that the corresponding Lagrangian is $L = \frac{1}{2}f(x)\dot{x}^2$.
- b) (25 p.) Write the transition amplitude in discrete space and integrate it over momentum variables.
- c) (15 p.) Take the limit $\Delta t \rightarrow 0$, $\Delta x = x_{i+1} - x_i \rightarrow 0$.
Hint: $\frac{1}{\Delta t}\delta_{ij} \rightarrow \delta(t_i - t_j)$.
- d) (5 p.) Show that

$$K(x_f, t_f; x_i, t_i) = \int \mathcal{D}x(t) \exp \left[\frac{i}{\hbar} S_{\text{eff}} \right],$$

with

$$S_{\text{eff}} = \int dt \left[L - \frac{i\hbar}{2} \delta(0) \ln f(x) \right].$$

Exercise 2. Transition amplitude of a free particle (50 points)

Use the expression for the transition amplitude $K_0(x, t; 0, 0)$ for a free particle in one dimension as derived in the lecture. Show that K_0 satisfies a Schroedinger equation:

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} K_0(x, t; 0, 0) = i\hbar \frac{\partial}{\partial t} K_0(x, t; 0, 0).$$