

PATH INTEGRALS


IN

QUANTUM MECHANICS

- 1) DERIVATION OF PATH INTEGRAL IN 1D
- 2) APPLICATION OF PATH INTEGRALS IN 1D
- 3) EXAMPLE IN 3D : AHARONOV-BOHM EFFECT
- 4) BROWNIAN MOTION AND WIENER PATH INTEGRAL

1) DERIVATION OF PATH INTEGRAL IN 1D

• CLASSICAL ACTION



$$\text{ACTION } S[x(t)] = \int_{t_i}^{t_f} dt \, L(x(t), \dot{x}(t))$$

CONSIDER VARIATIONS OF S BY $\delta x(t)$

SUCH THAT $\delta x(t_i) = 0$, $\delta x(t_f) = 0$

$x(t_i) = x_i$, $x(t_f) = x_f$

\Rightarrow VARIATIONAL PRINCIPLE (PRINCIPLE OF LEAST ACTION)

CLASSICAL PATH MINIMIZES S

$$\text{i.e. } \underline{\underline{\delta S = 0}}$$


\Downarrow
EULER-LAGRANGE EQ.

$$\underline{\underline{\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) - \frac{\partial L}{\partial x} = 0}}$$

\hookrightarrow SOLUTION GIVES CLASSICAL PATH


• TRANSITION AMPLITUDE IN QUANTUM MECHANICS

↳ CONSIDER 2-SLIT EXP.




AT POSITION x_f & TIME t_f
QM AMPLITUDE IS OBTAINED
AS SUM OF AMPLITUDE
FOR 2 PATHS
(INTERFERE)

↳ MAKE MORE SLITS



4 SLITS
QM AMPLITUDE AT x_f, t_f
OBTAINED AS SUM
OVER 4 PATHS

↳ MAKE INFINITE # SLITS \rightarrow INTERMEDIATE SCREEN DISAPPEARS



QM AMPLITUDE AT x_f, t_f
OBTAINED AS SUM
OVER ALL POSSIBLE PATHS

↳ INSERT COMPLETE SET OF STATES OF H

$$\begin{aligned}
 & K(x_f, t_f; x_i, t_i) \\
 &= \langle x_f, t_f | x_i, t_i \rangle \\
 &= \langle x_f | e^{-\frac{i}{\hbar} \hat{H} (t_f - t_i)} | x_i \rangle \\
 &= \sum_m \sum_{m'} \langle x_f | \Psi_m \rangle \langle \Psi_m | e^{-\frac{i}{\hbar} \hat{H} (t_f - t_i)} | \Psi_{m'} \rangle \langle \Psi_{m'} | x_i \rangle
 \end{aligned}$$

$|\Psi_m\rangle$: EIGENSTATES OF \hat{H}

$$\hat{H} |\Psi_m\rangle = E_m |\Psi_m\rangle$$

$$\langle \Psi_m | \Psi_{m'} \rangle = \delta_{mm'}$$

$$= \sum_m \underbrace{\langle x_f | \Psi_m \rangle}_{\Psi_m(x_f)} e^{-\frac{i}{\hbar} E_m (t_f - t_i)} \underbrace{\langle \Psi_m | x_i \rangle}_{\Psi_m^*(x_i)}$$

$$= \sum_m e^{-\frac{i}{\hbar} E_m (t_f - t_i)} \Psi_m(x_f) \Psi_m^*(x_i)$$

FOURIER ANALYZING GIVES EIGENVALUES E_m OF \hat{H}

FOURIER COEFFICIENTS EIGENSTATES $|\Psi_m\rangle$


PROPAGATOR CONTAINS ALL DYNAMICAL INFORMATION ON QUANTUM SYSTEM

• TRANSITION AMPLITUDE AS PATH INTEGRAL

R.P. FEYNMAN (1948)

↳ BREAK UP TIME INTERVAL $t_f - t_i$ INTO N INFINITESIMAL

INTERVALS Δt



$$t_k = t_i + k \Delta t$$

+ USE COMPLETENESS OF EIGENSTATES
AT INTERMEDIATE t_k

$$\int_{-\infty}^{+\infty} dx_k |x_k, t_k\rangle \langle x_k, t_k| = \mathbb{1}$$

↳ $K(x_f, t_f; x_i, t_i)$

$$= \prod_{k=1}^{N-1} \int dx_k \langle x_N, t_N | x_{N-1}, t_{N-1} \rangle \langle x_{N-1}, t_{N-1} | x_{N-2}, t_{N-2} \rangle \dots$$

$$\dots \langle x_2, t_2 | x_1, t_1 \rangle \langle x_1, t_1 | x_0, t_0 \rangle$$

↳ FOR INFINITESIMAL INTERVAL

$$\langle x_{k+1}, t_{k+1} | x_k, t_k \rangle$$

$$= \langle x_{k+1} | e^{-\frac{i}{\hbar} \hat{H}(t_{k+1} - t_k)} | x_k \rangle$$

$$= \langle x_{k+1} | e^{-\frac{i}{\hbar} \hat{H}(\Delta t)} | x_k \rangle$$

INSERT COMPLETE SET OF MOMENTUM
EIGENSTATES

$$\hat{P} |P_k\rangle = P_k |P_k\rangle$$

$$\langle P_k | P_{k'} \rangle = \delta(P_{k'} - P_k)$$

$$\int_{-\infty}^{+\infty} dP_k |P_k\rangle \langle P_k| = \mathbb{1}$$

$$\therefore \langle x_{k+1}, t_{k+1} | x_k, t_k \rangle$$

$$= \int dP_k \langle x_{k+1} | P_k \rangle \langle P_k | e^{-\frac{i}{\hbar} \hat{H} \Delta t} | x_k \rangle$$

$$\downarrow \quad \hat{H} \quad \text{DEPENDS ON } \hat{x} \text{ \& } \hat{P}$$

$$= \int dP_k \langle x_{k+1} | P_k \rangle \langle P_k | x_k \rangle e^{-\frac{i}{\hbar} H(x_k, P_k) \Delta t}$$

NOTE: THIS IS THE
HAMILTONIAN FUNCTION
NOT OPERATOR ANY MORE

$$\text{e.g. } H(x, p) = \frac{p^2}{2m} + V(x)$$

↳ USE

$$\int dP_k \langle x_{k+1} | P_k \rangle \langle P_k | x_k \rangle = \langle x_{k+1} | x_k \rangle$$

$$= \delta(x_{k+1} - x_k)$$

$$= \frac{1}{2\pi\hbar} \int dP_k e^{+\frac{i}{\hbar} P_k (x_{k+1} - x_k)}$$

$$\langle x_{k+1} | P_k \rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{\frac{i}{\hbar} P_k x_{k+1}}$$

$$\langle P_k | x_k \rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{-\frac{i}{\hbar} P_k x_k}$$

$$\hookrightarrow \langle x_{k+1}, t_{k+1} | x_k, t_k \rangle$$

$$= \int \frac{dP_k}{2\pi\hbar} e^{\frac{i}{\hbar} [P_k (x_{k+1} - x_k) - H(x_k, P_k) \cdot \Delta t]}$$

↓ FOR Δt INFINITESIMAL

$$x_{k+1} - x_k = \dot{x}_k (\Delta t)$$

$$\langle x_{k+1}, t_{k+1} | x_k, t_k \rangle = \int \frac{dP_k}{2\pi\hbar} e^{\frac{i}{\hbar} [P_k \dot{x}_k - H(x_k, P_k)] \Delta t}$$

↳ FINITE TRANSITION AMPLITUDE

$$\begin{aligned}
 & K(x_f, t_f; x_i, t_i) \\
 &= \prod_{k=1}^{N-1} \int dq_k \prod_{k=0}^{N-1} \int \frac{dp_k}{2\pi\hbar} \\
 &\quad \cdot \exp \left\{ \frac{i}{\hbar} \sum_{k=0}^{N-1} \left[p_k \dot{x}_k - H(x_k, p_k) \right] \Delta t \right\} \\
 &\quad \xrightarrow{\Delta t \rightarrow 0} \int_{t_i}^{t_f} dt \left[p \dot{x} - H(x, p) \right]
 \end{aligned}$$

↳ PATH INTEGRAL

$$\begin{aligned}
 & K(x_f, t_f; x_i, t_i) \\
 &= \int \mathcal{D}x(t) \mathcal{D}p(t) \exp \left\{ \frac{i}{\hbar} \int_{t_i}^{t_f} dt \left[p \dot{x} - H(x, p) \right] \right\}
 \end{aligned}$$

WITH PATH INTEGRAL 'MEASURES'

$$\mathcal{D}x(t) \equiv \lim_{N \rightarrow \infty} \prod_{k=1}^{N-1} dx(t_k)$$

$$\mathcal{D}p(t) \equiv \lim_{N \rightarrow \infty} \prod_{k=0}^{N-1} \frac{dp(t_k)}{2\pi\hbar}$$

WITH $x(t_i) = x_i$ AND $x(t_f) = x_f$

NOTE : IN EXPONENTIAL WE HAVE CLASSICAL
ACTION FUNCTION OF SYSTEM
 IN TERMS OF HAMILTONIAN

$$S = \int_{t_i}^{t_f} dt [p \dot{x} - H(x, p)]$$

∴ PATH INTEGRAL IS

→ FUNCTIONAL INTEGRAL OVER ALL POSSIBLE
 TRAJECTORIES IN PHASE SPACE OF SYSTEM

→ WEIGHTED BY $e^{\frac{i}{\hbar} S}$ WITH
 S THE HAMILTONIAN ACTION

PATH INTEGRAL IN TERMS OF LAGRANGIAN ACTION

FOR $H(x, p) = \frac{p^2}{2m} + V(x)$

WE CAN PERFORM THE p_k INTEGRATIONS (FORMALLY)

$$\hookrightarrow \langle x_{k+1}, t_{k+1} | x_k, t_k \rangle$$

$$= \int \frac{dp_k}{2\pi\hbar} e^{\frac{i}{\hbar} [p_k \dot{x}_k - H(x_k, p_k)] \Delta t}$$

$$= e^{-\frac{i}{\hbar} V(x_k) \Delta t} \int \frac{dp_k}{2\pi\hbar} e^{\frac{i}{\hbar} \left[-\frac{p_k^2}{2m} + p_k \dot{x}_k \right] \Delta t}$$

$$= e^{-\frac{i}{\hbar} V(x_k) \Delta t} e^{+\frac{i}{\hbar} \frac{1}{2} m \dot{x}_k^2} \int \frac{dp_k}{2\pi\hbar} e^{-\frac{i \Delta t}{\hbar 2m} (p_k - m \dot{x}_k)^2}$$

ANALYTICAL CONTINUATION TO IMAG. TIME

$$\Delta \tau = i \Delta t \quad \text{REAL}$$

+ GAUSSIAN INTEGRAL

$$\int_{-\infty}^{+\infty} dx e^{-ax^2} = \sqrt{\frac{\pi}{a}}$$

$$= e^{+\frac{i}{\hbar} \left[\frac{1}{2} m \dot{x}_k^2 - V(x_k) \right] \Delta t} \cdot \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2}$$

$$= \underline{\underline{\left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} e^{\frac{i}{\hbar} L(x_k, \dot{x}_k) \Delta t}}}$$

$$\begin{aligned}
 & \langle x_{k+1}, t_{k+1} | x_k, t_k \rangle \\
 &= \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} e^{\frac{i}{\hbar} L(x_k, \dot{x}_k) \Delta t}
 \end{aligned}$$

WITH LAGRANGIAN FUNCTION

$$L(x, \dot{x}) = \frac{1}{2} m \dot{x}^2 - V(x)$$

↳ FINITE TRANSITION AMPLITUDE

$$\begin{aligned}
 & K(x_f, t_f; x_i, t_i) \\
 &= \lim_{N \rightarrow \infty} \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{N/2} \int_{-\infty}^{+\infty} \prod_{k=1}^{N-1} dx_k e^{\frac{i}{\hbar} \sum_{k=0}^{N-1} L(x_k, \dot{x}_k) \Delta t}
 \end{aligned}$$

⇓

$$\begin{aligned}
 & K(x_f, t_f; x_i, t_i) \\
 &= \int \tilde{\mathcal{D}}_x(t) \exp \left\{ \frac{i}{\hbar} \int_{t_i}^{t_f} dt L(x, \dot{x}) \right\} \\
 &= \int \tilde{\mathcal{D}}_x(t) \exp \left\{ \frac{i}{\hbar} S \right\}
 \end{aligned}$$

WITH

$$\tilde{\mathcal{D}}_x(t) \equiv \lim_{N \rightarrow \infty} \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{N/2} \prod_{k=1}^{N-1} dx(t_k)$$

PATH INTEGRAL

↳ SUM OVER ALL PATHS $x(t)$

↳ EACH PATH WEIGHTED WITH $e^{\frac{i}{\hbar} S}$

↳ CLASSICAL LIMIT $\hbar \rightarrow 0$

$e^{\frac{i}{\hbar} S}$ OSCILLATES RAPIDLY WHEN $\hbar \rightarrow 0$

ONLY PATH WHICH MAKES ACTION S STATIONARY
CONTRIBUTES TO PATH INTEGRAL

$$\delta S[x] = 0$$



$$x(t) = x_{cl}(t)$$

CLASSICAL PATH
(SOLUTION OF EULER-LAGRANGE EQ.)

• EQUIVALENCE WITH SCHRÖDINGER EQUATION

↳ WAVEFUNCTION

$$\begin{aligned}
 \Psi(x_f, t_f) &= \langle x_f, t_f | \Psi(t=0) \rangle \\
 &= \langle x_f | \underbrace{e^{-\frac{i}{\hbar} \hat{H} t_f}}_{| \Psi(t_f) \rangle} | \Psi(t=0) \rangle \\
 &= \langle x_f | \Psi(t_f) \rangle
 \end{aligned}$$

↳ CONNECTION WITH TRANSITION AMPLITUDE

$$\begin{aligned}
 \Psi(x_f, t_f) &= \int dx_i \langle x_f, t_f | x_i, t_i \rangle \langle x_i, t_i | \Psi(t=0) \rangle \\
 &= \int dx_i K(x_f, t_f; x_i, t_i) \Psi(x_i, t_i)
 \end{aligned}$$

↳ CONSIDER INFINITESIMAL TIME STEP

$$\begin{aligned}
 t_i &= t \\
 t_f &= t + \Delta t \quad (\Delta t \rightarrow 0)
 \end{aligned}$$

$$K(x_f, t + \Delta t; x_i, t) = \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} e^{\frac{i}{\hbar} L\left(\frac{x_i + x_f}{2}, \frac{x_f - x_i}{\Delta t}\right) \Delta t}$$

$$\Psi(x_f, t + \Delta t) = \int dx_i K(x_f, t + \Delta t; x_i, t) \Psi(x_i, t)$$

↳ CONSIDER PARTICLE MOVING IN 1D

$$L = \frac{1}{2} m \dot{x}^2 - V(x, t)$$

$$\rightsquigarrow K(x_f, t + \Delta t; x_i, t) = \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} e^{\frac{i}{\hbar} \frac{m}{2\Delta t} (x_f - x_i)^2 - \frac{i}{\hbar} \Delta t V\left(\frac{x_i + x_f}{2}, t\right)}$$

$x_f \equiv x$
 $x_i \equiv y$

$$\rightsquigarrow \Psi(x, t + \Delta t) = \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} \int_{-\infty}^{+\infty} dy e^{-\frac{m}{2\hbar i \Delta t} (x-y)^2 - \frac{i}{\hbar} \Delta t V\left(\frac{x+y}{2}, t\right)} \Psi(y, t)$$

ONLY REGION $y \approx x$ CONTRIBUTES MAINLY TO INTEGRAL
(ONLY REGION AROUND $\eta \approx 0$)

$$y = x + \eta$$

CHANGE INTEGRATION VARIABLE

$$\Psi(x, t + \Delta t) = \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} \int_{-\infty}^{+\infty} d\eta e^{-\frac{m \eta^2}{2\hbar i \Delta t} - \frac{i}{\hbar} \Delta t V\left(x + \frac{\eta}{2}, t\right)} \Psi(x + \eta, t)$$

EXPAND IN Δt AND KEEP ONLY LINEAR TERMS IN Δt & UP TO QUADRATIC TERMS IN η

$$\begin{aligned} & \Psi(x, t) + \Delta t \frac{\partial \Psi}{\partial t} \\ &= \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} \int_{-\infty}^{+\infty} d\eta e^{-\frac{m}{2\hbar i \Delta t} \eta^2} \left[1 - \frac{i}{\hbar} \Delta t V(x, t) + \dots \right] \\ & \quad \cdot \left[\Psi(x, t) + \eta \frac{\partial \Psi}{\partial x} + \frac{1}{2} \eta^2 \frac{\partial^2 \Psi}{\partial x^2} + \dots \right] \end{aligned}$$

$$\rightsquigarrow \left(\frac{a}{\pi}\right)^{1/2} \int_{-\infty}^{+\infty} d\eta e^{-a\eta^2} = 1 \quad \left(a = \frac{m}{2\hbar i \Delta t}\right)$$

$$\left(\frac{a}{\pi}\right)^{1/2} \int_{-\infty}^{+\infty} d\eta \eta e^{-a\eta^2} = 0$$

$$\begin{aligned} \left(\frac{a}{\pi}\right)^{1/2} \int_{-\infty}^{+\infty} d\eta \eta^2 e^{-a\eta^2} &= \left(\frac{a}{\pi}\right)^{1/2} \left(-\frac{1}{2a}\right) \int_{-\infty}^{+\infty} d(e^{-a\eta^2}) \eta \\ &= \left(\frac{a}{\pi}\right)^{1/2} \frac{1}{2a} \int_{-\infty}^{+\infty} d\eta e^{-a\eta^2} \\ &= \frac{1}{2a} \end{aligned}$$

$$\begin{aligned} \rightsquigarrow \psi(x,t) + \Delta t \frac{\partial \psi}{\partial t} \\ = \psi(x,t) - \frac{i}{\hbar} \Delta t V(x,t) \psi(x,t) \\ + \left(\frac{\hbar i \Delta t}{m}\right) \cdot \frac{1}{2} \frac{\partial^2 \psi}{\partial x^2} \end{aligned}$$

IDENTIFY TERM $O(\Delta t)$

$$\frac{\partial \psi}{\partial t} = -\frac{i}{\hbar} \left\{ -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V(x,t) \psi \right\}$$

↓
SCHRÖDINGER EQ !

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V(x,t) \psi = i\hbar \frac{\partial \psi}{\partial t}$$

2) APPLICATION OF PATH INTEGRALS IN 1D

- FREE PARTICLE ($V=0$)

$$L(x, \dot{x}) = \frac{1}{2} m \dot{x}^2$$

$$\rightsquigarrow K_0(x_f, t_f; x_i, t_i) = \int_{x_i}^{x_f} \tilde{\mathcal{D}}x(t) e^{\frac{i}{\hbar} \int_{t_i}^{t_f} dt \left(\frac{1}{2} m \dot{x}^2 \right)}$$

K_0

0 DENOTES
FREE PARTICLE

$$= \lim_{N \rightarrow \infty} \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{N/2} \int_{-\infty}^{+\infty} \prod_{k=1}^{N-1} dx_k e^{\frac{i}{\hbar} \Delta t \sum_{k=0}^{N-1} \frac{1}{2} m \frac{(x_{k+1} - x_k)^2}{(\Delta t)^2}}$$

$$\text{WITH } \begin{cases} x_0 \equiv x_i \\ x_N \equiv x_f \end{cases}$$

\rightsquigarrow PERFORM FIRST $\int dx_1$

$$\left(\frac{m}{2\pi\hbar i \Delta t} \right)^{2/2} \int dx_1 e^{-\left(\frac{m}{2\hbar i \Delta t} \right) \left\{ (x_2 - x_1)^2 + (x_1 - x_0)^2 \right\}}$$

$$\begin{aligned} & (x_2 - x_1)^2 + (x_1 - x_0)^2 \\ &= 2x_1^2 - 2(x_0 + x_2)x_1 + x_0^2 + x_2^2 \\ &= 2 \left(x_1 - \frac{1}{2}(x_0 + x_2) \right)^2 + \frac{1}{2}(x_0 - x_2)^2 \end{aligned}$$

$$= \left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} \cdot \frac{1}{\sqrt{2}} e^{-\frac{1}{2} \left(\frac{m}{2\hbar i \Delta t} \right) (x_0 - x_2)^2}$$

∴ AFTER $\int dx_1$ INTEGRATION

$$\left(\frac{m}{2\pi\hbar i \underline{2\Delta t}} \right)^{1/2} e^{-\left(\frac{m}{2\hbar i \underline{2\Delta t}} \right) (x_2 - x_0)^2}$$

→ MULTIPLY WITH $\left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2}$ AND $\int dx_2 e^{-\frac{m}{2\hbar i \Delta t} (x_3 - x_2)^2}$

$$\left(\frac{m}{2\pi\hbar i \Delta t} \right)^{1/2} \cdot \left(\frac{m}{2\pi\hbar i \underline{2\Delta t}} \right)^{1/2}$$

$$\cdot \int dx_2 e^{-\frac{m}{2\hbar i \Delta t} \left\{ (x_3 - x_2)^2 + \frac{1}{2} (x_2 - x_0)^2 \right\}}$$

$$\begin{aligned} & \left(x_3 - x_2 \right)^2 + \frac{1}{2} \left(x_2 - x_0 \right)^2 \\ &= \frac{3}{2} x_2^2 - (2x_3 + x_0) x_2 + x_3^2 + \frac{1}{2} x_0^2 \\ &= \frac{3}{2} \left(x_2 - \frac{1}{3} (2x_3 + x_0) \right)^2 \\ &+ x_3^2 + \frac{1}{2} x_0^2 - \frac{1}{6} (4x_3^2 + 4x_0 x_3 + x_0^2) \\ &= \frac{1}{3} (x_3 - x_0)^2 \end{aligned}$$

$$= \left(\frac{m}{2\pi\hbar i \underline{3\Delta t}} \right)^{1/2} e^{-\left(\frac{m}{2\hbar i \underline{3\Delta t}} \right) (x_3 - x_0)^2}$$

→ AFTER $N-1$ INTEGRATIONS.

$$\left(\frac{m}{2\pi\hbar i N\Delta t} \right)^{1/2} e^{-\left(\frac{m}{2\hbar i N\Delta t} \right) \left(\begin{matrix} x_f & - & x_i \\ \parallel & & \parallel \\ x_N & & x_0 \end{matrix} \right)^2}$$

$$\begin{aligned} \circ \circ \quad K_0(x_f, t_f; x_i, t_i) \\ = \lim_{N \rightarrow \infty} \left(\frac{m}{2\pi\hbar i N \Delta t} \right)^{1/2} e^{-\frac{m}{2\hbar i N \Delta t} (x_f - x_i)^2} \end{aligned}$$

$$\downarrow \quad N \Delta t = t_f - t_i$$

$$\begin{aligned} K_0(x_f, t_f; x_i, t_i) \\ = \left(\frac{m}{2\pi\hbar i (t_f - t_i)} \right)^{1/2} \exp \left\{ \frac{i m}{2\hbar} \frac{(x_f - x_i)^2}{(t_f - t_i)} \right\} \end{aligned}$$

→ NOTE : FOR CLASSICAL FREE PARTICLE

$$x_{cl}(t) = x_i + \left(\frac{x_f - x_i}{t_f - t_i} \right) (t - t_i)$$

$$\dot{x}_{cl} = \frac{x_f - x_i}{t_f - t_i} \quad \text{CONSTANT}$$

$$S_{cl} = \int_{t_i}^{t_f} dt \left(\frac{1}{2} m \dot{x}_{cl}^2 \right)$$

$$S_{cl} = \frac{m}{2} \frac{(x_f - x_i)^2}{(t_f - t_i)}$$

FOR QUANTUM FREE PARTICLE

$$K_0(x_f, t_f; x_i, t_i) \sim e^{\frac{i}{\hbar} S_{cl}}$$

→ MOMENTUM OF FREE PARTICLE

TAKE FREE PARTICLE INITIALLY AT $x_i = 0, t_i = 0$

• CLASSICAL $S_{cl} = \frac{m}{2} \frac{x_f^2}{t_f}$

$\frac{\partial S_{cl}}{\partial x_f} = m \left(\frac{x_f}{t_f} \right) = P$
 ↑ VELOCITY v ↑ MOMENTUM

• QM : PROBABILITY AMPLITUDE FOR PARTICLE TO BE FOUND AT x_f, t_f

$K_0(x_f, t_f; 0, 0)$
 $= \left(\frac{m}{2\pi\hbar i t_f} \right)^{1/2} e^{i \frac{c}{\hbar} \frac{m}{2} \frac{x_f^2}{t_f}}$

↳ FOR FIXED t_f AS x_f VARIES
 OSCILLATES

⇓
 PARTICLE BEHAVES AS WAVE

WAVELENGTH : COMPUTED FROM PERIODICITY CONDITION

$2\pi = \frac{m}{2\hbar t_f} [(x_f + \lambda)^2 - x_f^2]$

↓ $x_f \gg \lambda$

$2\pi = \frac{m}{\hbar} \left(\frac{x_f}{t_f} \right) \lambda \Rightarrow \lambda = \frac{h}{P}$
 DE BROGLIE

WITH $P = m \frac{x_f}{t_f}$
 ↑
 CLASSICAL MOMENTUM

~> ENERGY OF FREE PARTICLE

- CLASSICAL

$$- \frac{\partial S_{cl}}{\partial t_f} = \frac{1}{2} m \frac{x_f^2}{t_f^2} = \frac{1}{2} m v^2 = E$$

- QM

FOR FIXED x_f

K_0 OSCILLATES AS t_f VARIES

FREQUENCY $\omega = \frac{2\pi}{T}$ T: PERIOD

$$2\pi = \frac{m}{2\hbar} x_f^2 \left[\frac{1}{t_f} - \frac{1}{t_f + T} \right]$$

$$= \frac{m x_f^2}{2\hbar} \frac{T}{(t_f + T) t_f}$$

$$\downarrow \quad t_f \gg T$$


$$\frac{2\pi}{T} \approx \frac{m}{2\hbar} \frac{x_f^2}{t_f} = \frac{E}{\hbar}$$

$$\boxed{E = \hbar \omega}$$

CLASSICAL ENERGY $E = \frac{1}{2} m \frac{x_f^2}{t_f^2}$

SPREADING OF WAVEPACKET

GIVEN $\Psi(x_i, t=0)$ e.g.



HOW WILL $\Psi(x_f, t)$ LOOK LIKE?

SOLUTION THROUGH PATH INTEGRAL

(*)
$$\Psi(x_f, t) = \int dx_i K_0(x_f, t; x_i, 0) \Psi(x_i, 0)$$

$$K_0(x_f, t; x_i, 0) = \left(\frac{m}{2\pi\hbar i t} \right)^{1/2} e^{i \frac{m}{\hbar} \frac{(x_f - x_i)^2}{2t}}$$

⇓
USE FOURIER TF. TO EXPRESS INITIAL W.F. AS

$$\Psi(x_i, 0) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk e^{ikx_i} \Phi(k)$$

↓ USING (*)

$$\Psi(x_f, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk \Phi(k) \left(\frac{m}{2\pi\hbar i t} \right)^{1/2} \int dx_i e^{i \frac{m}{\hbar} \frac{(x_f - x_i)^2}{2t}} e^{ikx_i}$$

↓ CHANGE VARIABLE $\tilde{x}_i = x_i - x_f$

$$\psi(x_f, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk \phi(k) e^{ikx_f}$$

$$\left(\frac{m}{2\pi\hbar it}\right)^{1/2} \int d\tilde{x}_i e^{\underbrace{\left(\frac{i m \tilde{x}_i^2}{\hbar 2t} + ik\tilde{x}_i\right)}_{+\frac{i m}{\hbar 2t} \left(\tilde{x}_i + \frac{\hbar k}{m}\right)^2 - \frac{\hbar^2 k^2 t}{2m}}}$$

$$\downarrow \left(\frac{m}{2\pi\hbar it}\right)^{1/2} \int d\tilde{x}_i e^{-\frac{m}{\hbar 2it} \left(\tilde{x}_i + \frac{\hbar k}{m}\right)^2} = 1$$

$$\psi(x_f, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dk \phi(k) e^{\underbrace{-\frac{i}{\hbar} \left(\frac{\hbar^2 k^2}{2m}\right) t + ikx}_{\text{TRAVELLING WAVE}}}$$

TRAVELLING WAVE

→
WITH ENERGY $E = \frac{\hbar^2 k^2}{2m}$

MOMENTUM $\hbar k$

$\psi(x_f, t)$ DESCRIBES TIME
DEPENDENCE OF WAVE PACKET

HARMONIC OSCILLATOR

$$L(x, \dot{x}) = \frac{1}{2} m \dot{x}^2 - \frac{1}{2} m \omega^2 x^2$$

$$\rightsquigarrow K(x, t; x_i, 0) = \int_{x_i}^{x(t)=x} \tilde{\mathcal{D}} x(t') e^{\frac{i}{\hbar} S[x(t')]}$$

$$(\text{CHOOSE } t_i = 0) \quad x(0) = x_i$$

$$S[x(t')] = \int_0^t dt' \left[\frac{1}{2} m \dot{x}^2 - \frac{1}{2} m \omega^2 x^2 \right]$$

CLASSICAL PATH

$x_{cl}(t)$ IS SOLUTION OF E.L. EQ.

$$\frac{\partial L}{\partial x} - \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial \dot{x}} \right) = 0$$

$$-m\omega^2 x - m\ddot{x} = 0$$

$$\underline{\underline{\ddot{x} = -\omega^2 x}}$$

$$\downarrow$$

$$x_{cl}(t') = A \cos \omega t' + B \sin \omega t'$$

CONSTRAINTS $x_{cl}(0) = x_i \rightarrow A = x_i$

$$x_{cl}(t) = x \rightarrow B = \frac{x - x_i \cos \omega t}{\sin \omega t}$$

$$x_{cl}(t') = x_i \cos \omega t' + \frac{(x - x_i \cos \omega t)}{\sin \omega t} \sin \omega t'$$

→ CLASSICAL ACTION

$$\begin{aligned}
 S_{cl} \equiv S[x_{cl}] &= \frac{m\omega^2}{2} \int_0^t dt' \left\{ \left[-A \sin \omega t' + B \cos \omega t' \right]^2 \right. \\
 &\quad \left. - \left[A \cos \omega t' + B \sin \omega t' \right]^2 \right\} \\
 &= \frac{m\omega^2}{2} \int_0^t dt' \left\{ + \cos 2\omega t' (B^2 - A^2) \right. \\
 &\quad \left. - \sin 2\omega t' 2AB \right\} \\
 &= \frac{m\omega}{4} \left\{ + (B^2 - A^2) \sin 2\omega t' \Big|_0^t \right. \\
 &\quad \left. + 2AB \cos 2\omega t' \Big|_0^t \right\} \\
 &= \frac{m\omega}{4} \left\{ (-A^2 + B^2) \sin 2\omega t + 2AB (\cos 2\omega t - 1) \right\}
 \end{aligned}$$

$$S[x_{cl}] = \frac{m\omega}{2 \sin \omega t} \left\{ (x^2 + x_i^2) \cos \omega t - 2x x_i \right\}$$

→ QUANTUM ACTION

$$x(t') = x_{cl}(t') + Y(t)$$

↑
DEVIATION FROM CLASSICAL PATH

$$Y(0) = 0$$

$$Y(t) = 0$$

$$S[x(t')] = \frac{m}{2} \int_0^t dt' \left[(\dot{x}_{cl} + \dot{Y})^2 - \omega^2 (x_{cl} + Y)^2 \right]$$

$$= \frac{m}{2} \int_0^t dt' \left\{ \dot{x}_{cl}^2 - \omega^2 x_{cl}^2 + 2\dot{x}_{cl}\dot{Y} - 2\omega^2 x_{cl}Y + \dot{Y}^2 - \omega^2 Y^2 \right\}$$

NOTE : $\int_0^t dt' 2\dot{x}_{cl}\dot{Y} = 2\dot{x}_{cl}Y \Big|_0^t - \int_0^t dt' 2Y\ddot{x}_{cl}$

BECAUSE $Y(t) = Y(0) = 0$

$$\hookrightarrow \int_0^t dt' (2\dot{x}_{cl}\dot{Y} - 2\omega^2 x_{cl}Y)$$

$$= -2 \int_0^t dt' Y (\ddot{x}_{cl} + \omega^2 x_{cl})$$

$$= 0$$

↙ CLASSICAL EQUATION OF MOTION

$$\circ \circ \quad \left\| \begin{aligned} S[x(t')] &= S[x_{cl}] \\ &+ \frac{m}{2} \int_0^t dt' [\dot{y}^2 - \omega^2 y^2] \end{aligned} \right.$$

NOTE THAT 2nd TERM DOES NOT
DEPEND ON x OR x_i

WE CAN THEREFORE WRITE:

$$K(x, t; x_i, 0) = K(0, t; 0, 0) e^{\frac{i}{\hbar} S[x_{cl}]}$$

→ SHOW THAT

$$K(0, t; 0, 0) = \left(\frac{m\omega}{2\pi i\hbar \sin\omega t} \right)^{1/2}$$

$$\rightarrow K(x, t; x_i, 0)$$

$$= \left(\frac{m\omega}{2\pi i\hbar \sin\omega t} \right)^{1/2} e^{\frac{i}{\hbar} S[x_{cl}]}$$

→ NOTE : IN GENERAL → WHEN S CAN BE EXPRESSED
THROUGH A QUADRATIC FORM

$$K \sim e^{\frac{i}{\hbar} S[x_{cl}]}$$


Let's show that

$$K(0, t, 0, 0) = \left(\frac{m \omega}{2\pi i \hbar \sin(\omega t)} \right)^{1/2}$$

Due to our replacement

$$X(t') = X_{cl}(t') + y(t')$$

deviation from classical path



$$K(0, t, 0, 0) = \int_0^0 \mathcal{D}y(t) \exp\left(\frac{i}{\hbar} \frac{m}{2} \int_0^t dt' (\dot{y}^2 - \omega^2 y)\right)$$

such path can be written as a Fourier sine series with a fundamental period of t

$$y(t') = \sum_{n=1}^{\infty} A_n \sin\left(\frac{n\pi t'}{t}\right)$$

and it is possible to specify a path through the coefficients A_n instead of the function values $y(t_k)$.

The jacobian of this transformation J doesn't depend on ω .

General: all the prefactors that do not depend on ω we will recover from $\omega=0$ limit which corresponds to free particle

$$K(0, t, 0, 0) \xrightarrow{\omega=0} \left(\frac{m}{2\pi i \hbar t} \right)^{1/2}$$

Plug in $y(t')$ into $\exp(-)$:

$$\begin{aligned} 1) \quad \frac{m}{2} \int_0^t dt' \dot{y}^2 &= \frac{m}{2} \int_0^t dt' \sum_{n=1}^{\infty} a_n \left(\frac{n\pi}{t}\right) \cos\left(\frac{n\pi t'}{t}\right) \sum_{m=1}^{\infty} a_m \left(\frac{m\pi}{t}\right) \cos\left(\frac{m\pi t'}{t}\right) \\ &= \frac{m}{2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \left(\frac{n\pi}{t}\right) \left(\frac{m\pi}{t}\right) \int_0^t dt' \cos\left(\frac{n\pi t'}{t}\right) \cos\left(\frac{m\pi t'}{t}\right) \end{aligned}$$

using the relation

$$\cos(a) \cos(b) = \frac{1}{2} (\cos(a-b) + \cos(a+b))$$

$$= \frac{m}{2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \left(\frac{n\pi}{t}\right) \left(\frac{m\pi}{t}\right) \frac{t}{2\pi} \left(\underbrace{\frac{\sin(\pi(n-m))}{n-m}}_{n \neq m} + \underbrace{\frac{\sin(\pi(n+m))}{n+m}}_{n \neq -m} \right)$$

$$= \frac{m}{2} \frac{t}{2} \sum_{n=1}^{\infty} \left(\frac{n\pi}{t}\right)^2 a_n^2 \quad \begin{array}{ll} n \neq m & 0 \\ n = m & \pi \end{array} \quad \begin{array}{ll} n \neq -m & 0 \\ n = -m & \pi \end{array}$$

not possible
 $n, m = \{1, \infty\}$

Similarly

using the relation

$$\sin(a) \sin(b) = \frac{1}{2} (\cos(a-b) - \cos(a+b))$$

$$2) \quad \frac{m\omega^2}{2} \int_0^t dt' y^2 = \frac{m\omega^2}{2} \frac{t}{2} \sum_{n=1}^{\infty} a_n^2$$

On the assumption that $[0, t]$ region is divided into discrete steps, there is only a finite number N of coefficients a_n

$$k(0, t, 0, 0) = (J \dots) \int_{-p}^{+p} da_1 \int_{-p}^{+p} da_2 \dots \int_{-p}^{+p} da_N \exp \left\{ \frac{i m t}{2} \sum_{n=1}^{\infty} \left(\left(\frac{n\pi}{t}\right)^2 - \omega^2 \right) a_n^2 \right\}$$

some factors that do not depend on ω

Since the exp. can be separated into factors, the integral over each coefficients a_n can be done separately

$$\int_{-\infty}^{+\infty} da_n \exp \left\{ \frac{im}{2\hbar} \frac{t}{z} \left(\frac{\pi^2 \hbar^2}{t^2} - \omega^2 \right) a_n^2 \right\} = (\dots) \cdot \left(\frac{\pi^2 \hbar^2}{t^2} - \omega^2 \right)^{-1/2} =$$

↑ doesn't depend on ω

Gaussian integral $\int_{-\infty}^{+\infty} dx e^{-ax^2} = \sqrt{\frac{\pi}{a}}$

$$= (\dots) \left(1 - \frac{\omega^2 t^2}{\pi^2 \hbar^2} \right)^{-1/2}$$

Therefore

$$k(0, t, 0, 0) = (\dots) \prod_{n=1}^N \left(1 - \frac{\omega^2 t^2}{\pi^2 \hbar^2} \right)^{-1/2} = (\dots) \left(\frac{\sin \omega t}{\omega t} \right)^{-1/2}$$

$N \rightarrow \infty$

Since

$$k(0, t, 0, 0) \xrightarrow{\omega=0} \left(\frac{m}{2\pi \hbar i t} \right)^{1/2}$$

$$\Rightarrow \boxed{k(0, t, 0, 0) = \left(\frac{m \omega}{2\pi i \hbar \sin(\omega t)} \right)^{1/2}} \quad \neq$$

• PROJECTION OF THE GROUND STATE:

FEYNMAN-KAC FORMULA

$$\begin{aligned}
 &\rightsquigarrow K(x, t; x', 0) \\
 &= \langle x, t | x', 0 \rangle \\
 &= \langle x | e^{-\frac{i}{\hbar} \hat{H} t} | x' \rangle \\
 &= \sum_m \langle x | \psi_m \rangle \langle \psi_m | x' \rangle e^{-\frac{i}{\hbar} E_m t}
 \end{aligned}$$

\rightsquigarrow FOR $x = x'$

$$\begin{aligned}
 &K(x, t; x, 0) \\
 &= \sum_m |\psi_m(x)|^2 e^{-\frac{i}{\hbar} E_m t}
 \end{aligned}$$

\Downarrow

$$\begin{aligned}
 &\int dx K(x, t; x, 0) \\
 &= \int dx \langle x | e^{-\frac{i}{\hbar} \hat{H} t} | x \rangle \\
 &= \text{Tr} e^{-\frac{i}{\hbar} \hat{H} t} \\
 &= \sum_m \underbrace{\int dx |\psi_m(x)|^2}_1 e^{-\frac{i}{\hbar} E_m t}
 \end{aligned}$$

$$\int dx K(x, t; x, 0) = \sum_n e^{-\frac{i}{\hbar} E_n t} = \text{Tr} e^{-\frac{i}{\hbar} \hat{H} t} = \sum_n \langle n | e^{-\frac{i}{\hbar} \hat{H} t} | n \rangle$$

ANALYTIC CONTINUATION TO IMAGINARY TIME

$$\beta \equiv \frac{it}{\hbar}$$

$$\int dx K(x, t; x, 0) = \sum_n e^{-\beta E_n} = \text{Tr} e^{-\beta \hat{H}}$$

"PARTITION FUNCTION"

FOR $\beta \in \mathbb{R}$ AND POSITIVE THIS CORRESPONDS TO STATISTICAL MECHANICS PROBLEM (β IS $\frac{1}{k_B T}$ WITH T : TEMPERATURE)

IN LIMIT $\beta \rightarrow \infty$ (ZERO TEMPERATURE LIMIT) ONLY GROUND STATE CONTRIBUTES TO SUM

$$\int dx K(x, -it/\hbar; x, 0) \xrightarrow{\beta \rightarrow \infty} e^{-\beta E_0}$$

GROUND STATE ENERGY (FEYNMAN - KAC FORMULA)

$$E_0 = \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \ln \int dx K(x, -it/\hbar; x, 0) = \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \ln \text{Tr} e^{-\beta \hat{H}}$$

→ EXAMPLE : HARMONIC OSCILLATOR.

$$\hookrightarrow K(x, t; x', 0)$$

$$= \left(\frac{m\omega}{2\pi i \hbar \sin \omega t} \right)^{1/2} e^{\frac{i}{\hbar} S_{cl}}$$

$$\text{WITH } S_{cl} = \frac{m\omega}{2 \sin \omega t} \left\{ (x^2 + x'^2) \cos \omega t - 2xx' \right\}$$

$$\hookrightarrow \int dx K(x, t; x, 0)$$

$$= \left(\frac{m\omega}{2\pi i \hbar \sin \omega t} \right)^{1/2} \int dx e^{-a x^2}$$

$$\downarrow \text{ WITH } a = -\frac{i}{\hbar} \frac{m\omega}{\sin \omega t} (\cos \omega t - 1)$$

$$= \frac{i}{\hbar} (m\omega) \frac{\sin(\omega t/2)}{\cos(\omega t/2)}$$

$$= \left(\frac{m\omega}{2\pi i \hbar \sin \omega t} \right)^{1/2} \left(\frac{\pi \hbar \cos(\omega t/2)}{i m \omega \sin(\omega t/2)} \right)^{1/2}$$

$$2 \sin \frac{\omega t}{2} \cos \frac{\omega t}{2}$$

$$= \frac{1}{2i} \frac{1}{\sin(\omega t/2)}$$

$$E_0 = \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \ln \int dx \cdot K(x, -i\hbar\beta; x, 0)$$

$$= \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \ln \left[\frac{1}{2i} \frac{1}{\sin\left(-i\frac{\hbar\omega}{2}\beta\right)} \right]$$

$$\downarrow \quad \frac{1}{\sin x} = \frac{2i e^{-ix}}{1 - e^{-2xi}}$$

$$= \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \ln \left[\frac{e^{-\beta\frac{\hbar\omega}{2}}}{1 - e^{-\beta\hbar\omega}} \right]$$

$$= \lim_{\beta \rightarrow \infty} \left(-\frac{1}{\beta}\right) \left\{ -\beta\frac{\hbar\omega}{2} + O\left(e^{-\beta\hbar\omega}\right) \right\}$$

$$\underline{\underline{E_0 = \frac{\hbar\omega}{2}}}$$

↳ IN GENERAL : PARTITION FUNCTION

$$\int dx \cdot K(x, -i\hbar\beta; x, 0) = \sum_n e^{-\beta E_n}$$

$$\frac{e^{-\beta\frac{\hbar\omega}{2}}}{1 - e^{-\beta\hbar\omega}} = e^{-\beta\frac{\hbar\omega}{2}} \sum_{n=0}^{\infty} e^{-\beta n\hbar\omega}$$

⇓

$$\underline{\underline{E_m = \hbar\omega \left(m + \frac{1}{2}\right)}}$$

→ GROUND STATE WAVE FUNCTION

$$K(x, -i\hbar\beta; x, 0) = \sum_m e^{-\beta E_m} |\psi_m(x)|^2$$

$$\xrightarrow{\beta \rightarrow \infty} e^{-\beta E_0} |\psi_0(x)|^2$$

H.O.: $K(x, -i\hbar\beta; x, 0)$

$$= \left(\frac{m\omega}{\pi\hbar 2i \sin(-i\hbar\omega\beta)} \right)^{1/2} \exp \left\{ \frac{-i}{\hbar} m\omega x^2 \tan\left(-i\frac{\hbar\omega\beta}{2}\right) \right\}$$

$$\frac{1}{2i \sin(-i\hbar\omega\beta)} = \frac{e^{-\hbar\omega\beta}}{1 - e^{-2\hbar\omega\beta}} \downarrow$$

$$i \tan\left(-i\frac{\hbar\omega\beta}{2}\right) = \frac{1 - e^{-\hbar\omega\beta}}{1 + e^{-\hbar\omega\beta}}$$

$$= \left(\frac{m\omega}{\pi\hbar} \right)^{1/2} \frac{e^{-\beta E_0}}{\left(1 - e^{-\beta 2\hbar\omega}\right)^{1/2}} \exp \left\{ -\frac{m\omega}{\hbar} x^2 \left[1 + O(e^{-\beta\hbar\omega}) \right] \right\}$$

$$\xrightarrow{\beta \rightarrow \infty} e^{-\beta E_0} \left(\frac{m\omega}{\pi\hbar} \right)^{1/2} \exp \left\{ -\frac{m\omega}{\hbar} x^2 \right\}$$

$$\psi_0(x) = \left(\frac{m\omega}{\pi\hbar} \right)^{1/4} \exp \left\{ -\frac{m\omega}{2\hbar} x^2 \right\}$$

GROUND STATE W.F. OF H.O.

3) EXAMPLE IN 3D: AHARONOV-BOHM EFFECT

- CONSIDER e^- MOVING IN MAGNETIC FIELD

→ HAMILTONIAN $H = \frac{1}{2m} (\bar{p} - \frac{e}{c} \bar{A})^2$
 (MINIMAL SUBSTITUTION)

→ LAGRANGE FUNCTION $L = \bar{p} \cdot \dot{\bar{q}} - H$
 \bar{q} : COORDINATE, $\dot{\bar{q}}$ VELOCITY
 \bar{p} : CONJUGATE MOMENTUM

$$\bar{p} = \frac{\partial L}{\partial \dot{\bar{q}}}$$

$$\dot{\bar{q}} = \frac{\partial H}{\partial \bar{p}} = \frac{1}{m} (\bar{p} - \frac{e}{c} \bar{A})$$

→ CANONICAL MOMENTUM $\bar{p} = \underbrace{m \dot{\bar{q}} + \frac{e}{c} \bar{A}}_{\text{KINETIC MOMENTUM}}$


→ $L = (m \dot{\bar{q}} + \frac{e}{c} \bar{A}) \cdot \dot{\bar{q}} - \frac{m}{2} \dot{\bar{q}}^2$

$$\underline{\underline{L = \frac{1}{2} m \dot{\bar{q}}^2 + \frac{e}{c} \dot{\bar{q}} \cdot \bar{A}}}$$


→ ACTION $S = \int_0^t dt' L(\bar{q}, \dot{\bar{q}})$

$$= \frac{m}{2} \int_0^t dt' \frac{d\bar{q}}{dt'} \cdot \frac{d\bar{q}}{dt'} + \frac{e}{c} \int_0^t dt' \frac{d\bar{q}}{dt'} \cdot \bar{A}$$

$$= \frac{m}{2} \int_{\text{PATH}} d\bar{q} \cdot \frac{d\bar{q}}{dt'} + \frac{e}{c} \int_{\text{PATH}} d\bar{q} \cdot \bar{A}$$



→ CONSIDER CONDUCTING RING
CURRENT ENTERS AT POINT 1, IS EXTRACTED AT POINT 2



CONDUCTING WIRE AROUND SOLENOID $\rightarrow \vec{B}$ -FIELD

→ PROBABILITY AMPLITUDE FOR ELECTRON
TO BE OBSERVED AT POINT 2

$$K(2, 1) \sim e^{\frac{i}{\hbar} S(\text{PATH 1})} + e^{\frac{i}{\hbar} S(\text{PATH 2})}$$

$$= e^{\frac{i}{\hbar} S(\text{PATH 1})} \left(1 + e^{\frac{i}{\hbar} \Delta S} \right)$$

$$\Delta S = S(\text{PATH 2}) - S(\text{PATH 1})$$

$$= \frac{e}{c} \left\{ \int_{\text{PATH 2}} d\vec{q} \cdot \vec{A} - \int_{\text{PATH 1}} d\vec{q} \cdot \vec{A} \right\}$$

$$= \frac{e}{c} \oint d\vec{q} \cdot \vec{A}$$

NOTE : KINETIC ENERGY TERMS ARE EQUAL
DUE TO SYMMETRY \rightarrow AND CANCEL IN DIFFERENCE

STOKES THEOREM

$$\Delta S = \frac{e}{c} \int d\vec{S} \cdot (\vec{\nabla} \times \vec{A})$$

$d\vec{S}$: NORMAL TO SURFACE OF CURRENT LOOP

$$= \frac{e}{c} \int d\vec{S} \cdot \vec{B}$$

$$\Delta S = \frac{e}{c} \Phi$$

↳ MAGNETIC FLUX

∴ PROB. AMPLITUDE $\sim 1 + e^{\frac{ie}{\hbar c} \Phi}$

$$\text{PROBABILITY} = |K(2,1)|^2$$

$e^{\frac{ie}{\hbar c} \Phi}$ TERM WILL GIVE INTERFERENCE PATTERN WHICH WILL CHANGE BY VARYING Φ (i.e. \vec{B})

CONSTRUCTIVE INTERFERENCE

$$\frac{e\Phi}{\hbar c} = 2\pi m \Rightarrow \Phi = \left(\frac{\hbar c}{e}\right) m = m \phi_0$$

$$m \in \mathbb{Z}$$

$$\phi_0 = \frac{\hbar c}{e}$$

QUANTUM OF FLUX

MAXIMUM CURRENT FOR Φ EQUAL TO AN INTEGER TIMES ϕ_0 : AHARONOV - BOHM EFFECT

↳ WAS FIRST OBSERVED EXPERIMENTALLY BY CHAMBERS (1960)

4) BROWNIAN MOTION AND WIENER PATH INTEGRAL


• RANDOM WALK IN 1 DIMENSION

↳ CONSIDER DISCRETE RANDOM WALK

DISCRETE TIME STEPS ϵ

DISCRETE SPATIAL STEPS l EITHER TO LEFT OR TO RIGHT

SUPPOSE : INITIALLY $t=0$, POSITION $x=0$



↳ PROBABILITY FOR LEFT STEP AND RIGHT STEP \Rightarrow EACH $\frac{1}{2}$
 SUCCESSIVE STEPS ARE STATISTICALLY INDEPENDENT
 (STOCHASTIC PROCESS)

PROBABILITY FOR TRANSITION
 FROM $x = jl$ TO $x = il$ DURING TIME ϵ
 $(i, j \in \mathbb{Z})$

$$\| P(il - jl, \epsilon) = \begin{cases} \frac{1}{2}, & |i - j| = 1 \\ 0, & \text{OTHERWISE} \end{cases}$$

- HOMOGENEOUS IN SPACE : ONLY DEPENDS ON $i - j$
- ISOTROPIC IN SPACE : SYMMETRIC UNDER $(i, j) \rightarrow (-i, -j)$

↳ DISCRETE RANDOM WALK : EXAMPLE OF MARKOV CHAIN

MARKOV CHAIN : CHARACTERIZED BY $(P(t_m), P(0))$

$P_{ij}(t_m)$: TRANSITION PROBABILITY FROM $j \rightarrow i$ AT TIME t_m

$P_i(0)$: INITIAL PROBABILITY DISTRIBUTION

$$P_i(t_m) = \sum_j P_{ij}(t_m) P_j(0)$$

NOTE $0 \leq P_i(0) \leq 1$ $\sum_i P_i(0) = 1$

$0 \leq P_{ij} \leq 1$ $\sum_i P_{ij} = 1$

|| TRANSITION PROB. TO STATE i AT TIME t_m
 DEPENDS ONLY ON STATE j AT TIME t_{m-1}
 AND NOT ON STATES AT EARLIER TIMES t_{m-2}, t_{m-3}, \dots

FOR DISCRETE RANDOM WALK.

$$P_{ij}(\epsilon) = P(i^l - j^l, \epsilon)$$

MARKOV CHAIN : SUCCESSIVE STEPS STATISTICALLY INDEPENDENT

CHOOSE $P_j(0) = \delta_{j0}$, i.e. $x(0) = 0$

↳ P IN MATRIX NOTATION

MATRIX ELEMENT ij

$$P(\epsilon) = \frac{1}{2} (R(\epsilon) + L(\epsilon))$$

$$R(\epsilon) : \text{STEP TO RIGHT} \quad (R(\epsilon))_{ij} = \delta_{i, j+1}$$

$$L(\epsilon) : \text{STEP TO LEFT} \quad (L(\epsilon))_{ij} = \delta_{i, j-1}$$

e.g.

$$\begin{aligned} P_i(\epsilon) &= \sum_j P_{ij}(\epsilon) P_j(0) \\ &= \frac{1}{2} \sum_j (R_{ij}(\epsilon) + L_{ij}(\epsilon)) \delta_{j0} \\ &= \frac{1}{2} (R_{i0}(\epsilon) + L_{i0}(\epsilon)) \\ &= \frac{1}{2} (\delta_{i1} + \delta_{i-1}) \end{aligned}$$

↳ AFTER m TIME STEPS

$$P_i(m\epsilon) = \sum_j (P^m(\epsilon))_{ij} \underbrace{P_j(0)}_{\delta_{j0}}$$

NOTE $P = \frac{1}{2} (R + L)$

$$P^m = \frac{1}{2^m} \sum_{k=0}^m \binom{m}{k} R^k L^{m-k}$$

(BINOMIAL FORMULA)

$$\text{AS } RL = LR = 1$$

$$\downarrow$$

$$P^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} R^{2k-n}$$

$$(P^n)_{ij} = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \underbrace{(R^{2k-n})}_{\delta_{i, j+2k-n}}_{ij}$$

$(P^n)_{ij} = \begin{cases} \frac{1}{2^n} \binom{n}{\frac{1}{2}(i-j+n)} & , \text{ if } i-j \leq n \\ & \text{ AND } i-j+n \text{ EVEN} \\ 0 & , \text{ OTHERWISE} \end{cases}$

$$\Downarrow$$

$$P_i(m\varepsilon) = \sum_j (P^m)_{ij} P_j(0)$$

$$= (P^m)_{i0}$$

$$= \begin{cases} \frac{1}{2^m} \binom{m}{\frac{1}{2}(i-m)} & , \text{ if } |i| \leq m \\ & \text{ AND } i-m \text{ EVEN} \\ 0 & , \text{ OTHERWISE} \end{cases}$$

NOTE

$$P(i\ell - j\ell, m\varepsilon) = (P^{(m\varepsilon)})_{ij}$$

PROPERTIES:

- 1) HOMOGENEOUS IN SPACE : DEPENDS ONLY ON $i - \ell$
- 2) " " TIME : DOES NOT DEPEND ON INITIAL TIME, ONLY ON TIME DIFFERENCE m
- 3) ISOTROPIC IN SPACE $P(-i\ell + j\ell, m\varepsilon) = P(i\ell - j\ell, m\varepsilon)$

↳ RECURSION FORMULA FOR BINOMIAL COEFF

$$\binom{m+1}{k} = \binom{m}{k} + \binom{m}{k-1}$$

PROOF:

$$\text{LEFT} = \frac{(m+1)!}{(m+1-k)! k!}$$

$$\text{RIGHT} = \frac{m!}{(m-k)! k!} + \frac{m!}{(m-k+1)! (k-1)!}$$

$$= \frac{m!}{(m+1-k)! k!} (m+1-k + k) = \frac{(m+1)!}{(m+1-k)! k!}$$

AS $P(i\ell - j\ell, n\epsilon) = \frac{1}{2^n} \binom{m}{k}$

FOR $k = \frac{1}{2} (i-j + m)$

(IF) $|i-j| \leq m$
 $i-j + m$ EVEN

$$P(i\ell - j\ell, (m+1)\epsilon)$$

$$= \frac{1}{2} \frac{1}{2^m} \binom{m}{k} + \frac{1}{2} \frac{1}{2^m} \binom{m}{k-1}$$

$$k = \frac{1}{2} (i-j + m+1)$$

$$= \frac{1}{2} (i-j+1 + m)$$

$$k-1 = \frac{1}{2} (i-j + m) - 1$$

$$= \frac{1}{2} (i-j-1 + m)$$

$$= \frac{1}{2} P((i-j+1)\ell, m\epsilon) + \frac{1}{2} P((i-j-1)\ell, m\epsilon)$$

OR FOR $x = (i-j)l$
 $t = n\varepsilon$

$$P(x, t+\varepsilon) = \frac{1}{2} P(x+l, t) + \frac{1}{2} P(x-l, t)$$

$$\frac{1}{\varepsilon} \left(P(x, t+\varepsilon) - P(x, t) \right)$$

$$= \frac{l^2}{2\varepsilon} \cdot \frac{1}{l^2} \left(P(x+l, t) - 2P(x, t) + P(x-l, t) \right)$$



MACROSCOPIC DESCRIPTION OF RANDOM WALK

LIMIT $\varepsilon \rightarrow 0$

$l \rightarrow 0$

$$\frac{l^2}{2\varepsilon} = D \quad \text{FINITE}$$

$$\frac{\partial}{\partial t} P(x, t) = D \frac{\partial^2}{\partial x^2} P(x, t)$$

DIFFUSION
EQUATION

D : DIFFUSION CONSTANT

↳ SOLUTION OF DIFFUSION EQ.

INITIAL CONDITION

$$t=0 \Rightarrow x=0 : P(x,0) = \delta(x)$$

$$\int dx P(x,0) = 1$$

$$P(x,t) = \int dk e^{ikx} \tilde{P}(k,t)$$

$$\tilde{P}(k,t) = \overset{\uparrow}{\int} \frac{dx}{2\pi} e^{-ikx} P(x,t)$$

$$\tilde{P}(k,0) = \frac{1}{2\pi}$$

$$\circ \circ \quad \frac{\partial}{\partial t} P(x,t) = D \frac{\partial^2}{\partial x^2} P(x,t)$$

↓

$$\frac{\partial}{\partial t} \tilde{P}(k,t) = -Dk^2 \tilde{P}(k,t)$$

$$\frac{\partial}{\partial t} \ln \tilde{P}(k,t) = -Dk^2$$

$$\Downarrow \int_0^t$$

$$\ln \frac{\tilde{P}(k,t)}{\tilde{P}(k,0)} = -Dk^2 t$$

$$\boxed{\tilde{P}(k,t) = \frac{1}{2\pi} \exp(-Dk^2 t)}$$

$$\begin{aligned}
 \rightsquigarrow P(x,t) &= \int_{-\infty}^{+\infty} dk e^{ikx} \frac{1}{2\pi} e^{-Dk^2 t} \\
 &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dk e^{-Dt \left(k - \frac{ix}{2Dt}\right)^2 - \frac{x^2}{4Dt}} \\
 &= \frac{1}{2\pi} \sqrt{\frac{\pi}{Dt}} e^{-\frac{x^2}{4Dt}}
 \end{aligned}$$

$$P(x,t) = \left(\frac{1}{4\pi Dt} \right)^{1/2} e^{-\frac{x^2}{4Dt}}$$

SATISFIES $\int dx P(x,t) = 1$

MEAN SQUARE DEVIATION

$$\begin{aligned}
 \langle x^2 \rangle_t &= \int dx x^2 P(x,t), \quad \text{NOTE } \langle x \rangle = 0 \\
 &= 2Dt
 \end{aligned}$$

$$\Downarrow \\
 \underline{\underline{\sqrt{\langle x^2 \rangle_t} = \sqrt{2Dt}}}$$

\hookrightarrow FOR ARBITRARY INITIAL CONDITION $x(t_0) = x_0$

$$\begin{aligned}
 &P(x,t; x_0, t_0) \\
 &= \left(\frac{1}{4\pi D(t-t_0)} \right)^{1/2} e^{-\frac{(x-x_0)^2}{4D(t-t_0)}}
 \end{aligned}$$

GAUSSIAN

PROBABILITY FOR RANDOM WALKER
WHICH INITIALLY STARTS AT x_0, t_0
TO BE FOUND AT A LATER TIME t AT x

• IMAGINARY TIME

↳ DIFFUSION EQ.

$$D \frac{\partial^2 P}{\partial x^2} = \frac{\partial P}{\partial t}$$

⇓ ANALYTICALLY CONTINUE TO
IMAGINARY TIME i.e. $t \Rightarrow it$

$$D \frac{\partial^2 P}{\partial x^2} = \frac{\partial P}{\partial(it)}$$

$$- D \frac{\partial^2 P}{\partial x^2} = i \frac{\partial P}{\partial t}$$

FOR $D = \frac{\hbar}{2m}$ THIS IS SCHRÖDINGER EQ. !


$$\begin{aligned} \text{↳ } P(x, it; x_0, it) \Big|_{D = \frac{\hbar}{2m}} \\ = \left(\frac{m}{2\pi\hbar i(t-t_0)} \right)^{1/2} \exp \left\{ + \frac{im}{2\hbar} \frac{(x-x_0)^2}{(t-t_0)} \right\} \end{aligned}$$

NOTE 1) THIS IS PRECISELY KERNEL $K(x, t; x_0, t)$
FOR A FREE PARTICLE IN QUANTUM MECHANICS !

- 2) DIFFUSION EQ : GAUSSIAN P (PROBABILITY)
SCHRÖDINGER EQ : OSCILLATORY KERNEL (PROB. AMPLITUDE!)
⇓
→ ADMITS WAVE SOLUTIONS.
→ INTERFERENCE EFFECTS

- 3) QM FREE PARTICLE SOLUTION MAY BE SEEN
AS ANALYTIC CONTINUATION TO IMAG. TIME OF INDETERMINISTIC
MOTION OF BROWNIAN PARTICLE

• WIENER PATH INTEGRAL (1921)



↳ CONSIDER BROWNIAN PARTICLE

PROBABILITY TO FIND BROWNIAN PARTICLE

WHICH STARTED AT $t=0$ AT IN $x=0$

AT TIME t_1 IN INTERVAL $[a_1, b_1]$

AT TIME t_2 IN INTERVAL $[a_2, b_2]$

AT TIME t_N IN INTERVAL $[a_N, b_N]$

PRODUCT OF PROBABILITIES DUE TO STOCHASTIC PROCESS

PROB $\{ x(t_1) \in [a_1, b_1], x(t_2) \in [a_2, b_2], \dots, x(t_N) \in [a_N, b_N] \}$


$$= \int_{a_1}^{b_1} dx_1 P(x_1, t_1; 0, 0) \int_{a_2}^{b_2} dx_2 P(x_2, t_2; x_1, t_1)$$

$$\dots \int_{a_N}^{b_N} dx_N P(x_N, t_N; x_{N-1}, t_{N-1})$$

WITH $P(x_{i+1}, t_{i+1}; x_i, t_i) = \left(\frac{1}{4\pi D(t_{i+1} - t_i)} \right)^{1/2} e^{-\frac{(x_{i+1} - x_i)^2}{4D(t_{i+1} - t_i)}}$

↳ CONSIDER $t_i - t_{i-1} = \Delta t_i \rightarrow 0$ (i.e. $N \rightarrow \infty$)
 $x_i - x_{i-1} = dx_i \rightarrow 0$

PROBABILITY THAT BROWNIAN PARTICLE
 MOVES ALONG A TRAJECTORY $x(t)$
 FROM $x=0$ AT $t=0$ TO $x(t_f)$.




$$P = \lim_{\substack{\Delta t_i \rightarrow 0 \\ N \rightarrow \infty}} \exp \left\{ - \sum_{i=1}^N \frac{(x_i - x_{i-1})^2}{4D(t_i - t_{i-1})} \right\} \prod_{i=1}^N \left(\frac{1}{4\pi D(t_i - t_{i-1})} \right)^{1/2} dx_i$$

$$= \lim_{\substack{\Delta t_i \rightarrow 0 \\ N \rightarrow \infty}} \exp \left\{ - \sum_{i=1}^N \left(\frac{x_i - x_{i-1}}{t_i - t_{i-1}} \right)^2 \frac{\Delta t_i}{4D} \right\} \prod_{i=1}^N \left(\frac{1}{4\pi D \Delta t_i} \right)^{1/2} dx_i$$

$$= \exp \left\{ - \frac{1}{4D} \int_0^{t_f} \dot{x}^2(t) dt \right\} \prod_{t=0}^{t_f} \frac{dx(t)}{\sqrt{4\pi D dt}}$$

↳ PATH INTEGRAL (WIENER)



$$P(x_f, t_f; 0, 0)$$

CAN BE OBTAINED IN 2 WAYS

1) AS BEFORE

$$P(x_f, t_f; 0, 0) = \left(\frac{1}{4\pi D t_f} \right)^{1/2} e^{-\frac{x_f^2}{4 D t_f}}$$

2) AS SUM OF PROBABILITIES OF ALL PATHS
CONNECTING $x(0) = 0$ AND $x(t_f) = x_f$

$$P(x_f, t_f; 0, 0) = \int_{C\{0,0; x_f, t_f\}} d_W x(t)$$

WITH $d_W x(t)$: WIENER MEASURE

$$d_W x(t) \equiv \exp \left\{ -\frac{1}{4D} \int_0^{t_f} dt \dot{x}^2(t) \right\} \prod_{t=0}^{t_f} \frac{dx(t)}{\sqrt{4\pi D dt}}$$

WELL DEFINED INTEGRAL

NOTE:

↳ FEYNMAN PATH INTEGRAL (FREE PARTICLE)

$$K(x, t_f ; 0, 0) \sim \exp \left\{ \frac{i}{\hbar} \int_0^{t_f} dt \frac{1}{2} m \dot{x}^2 \right\}$$

↓
ANALYTICALLY CONTINUE TO
IMAG TIME $\tau = it$

$$K(x, it_f ; 0, 0) \sim \exp \left\{ \frac{1}{\hbar} \int_0^{it_f} d(it) \frac{1}{2} m \dot{x}^2 \right\}$$

↓ $\dot{x}^2 = - \left(\frac{dx}{d\tau} \right)^2$

$$K(x, \tau ; 0, 0) \sim \exp \left\{ - \frac{m}{2\hbar} \int_0^{\tau_f} d\tau \left(\frac{dx}{d\tau} \right)^2 \right\}$$

$$= \exp \left\{ - \frac{1}{4D} \int_0^{\tau_f} d\tau \left(\frac{dx}{d\tau} \right)^2 \right\}$$

WITH $D = \frac{\hbar}{2m}$